


Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1




2344
2345
2346
2347
2348
2349
2350
2351
2352
2353
2354
2355
2356
2357
2358
2359
2360
2361

L. Burbank, M. C. Roper. OxyR and SoxR modulate the inducible oxidative stress response and are implicated during different stages of infection for the bacterial phytopathogen <i>Pantoea stewartii</i> subsp. <i>stewartii</i> . <i>Mol Plant Microbe Interact.</i> 2014. 27:479-90
M. Zheng, B. Doan, T. D. Schneider, G. Storz. OxyR and SoxRS regulation of <i>fur</i> . <i>J Bacteriol.</i> 1999. 181:4639-43
K. Xie, H. Peng, H. Hu, W. Wang, X. Zhang. OxyR, an important oxidative stress regulator to phenazines production and hydrogen peroxide resistance in <i>Pseudomonas chlororaphis</i> GP72. <i>Microbiol Res.</i> 2013. 168:646-53
S. I. Liochev, I. Fridovich. Paraquat diaphorases in <i>Escherichia coli</i> . <i>Free Radic Biol Med.</i> 1994. 16:555-9
J. Membrillo-Hernandez, S. O. Kim, G. M. Cook, R. K. Poole. Paraquat regulation of <i>hmp</i> (flavo-hemoglobin) gene expression in <i>Escherichia coli</i> K-12 is SoxRS independent but modulated by sigma S. <i>J Bacteriol.</i> 1997. 179:3164-70
Z. T. Wen, J. P. Bitoun, S. Liao. PBP1a-deficiency causes major defects in cell division, growth and biofilm formation by <i>Streptococcus mutans</i> . <i>PLoS One.</i> 2015. 10:e0124319
M. E. Achard, A. J. Hamilton, T. Dankowski, B. Heras, M. S. Schembri, J. L. Edwards, M. P. Jennings, A. G. McEwan. A periplasmic thioredoxin-like protein plays a role in defense against oxidative stress in <i>Neisseria gonorrhoeae</i> . <i>Infect Immun.</i> 2009. 77:4934-9
S. R. Palmer, J. H. Miller, J. Abranches, L. Zeng, T. Lefebure, V. P. Richards, J. A. Lemos, M. J. Stanhope, R. A. Burne. Phenotypic heterogeneity of genomically-diverse isolates of <i>Streptococcus mutans</i> . <i>PLoS One.</i> 2013. 8:e61358
K. Jantzer, K. Zerulla, J. Soppa. Phenotyping in the archaea: optimization of growth parameters and analysis of mutants of <i>Haloferax volcanii</i> . <i>FEMS Microbiol Lett.</i> 2011. 322:123-30
M. R. Engle, S. P. Singh, P. J. Czernik, D. Gaddy, D. C. Montague, J. D. Ceci, Y. Yang, S. Awasthi, Y. C. Awasthi, P. Zimniak. Physiological role of mGSTA4-4, a glutathione S-transferase metabolizing 4-hydroxynonenal: generation and analysis of mGsta4 null mouse. <i>Toxicol Appl Pharmacol.</i> 2004. 194:296-308
G. Mir, J. Domenech, G. Huguet, W. J. Guo, P. Goldsbrough, S. Atrian, M. Molinas. A plant type 2 metallothionein (MT) from cork tissue responds to oxidative stress. <i>J Exp Bot.</i> 2004. 55:2483-93
M. Hanikenne, R. F. Matagne, R. Loppes. Pleiotropic mutants hypersensitive to heavy metals and to oxidative stress in <i>Chlamydomonas reinhardtii</i> . <i>FEMS Microbiol Lett.</i> 2001. 196:107-11
E. J. Schott, G. R. Vasta. The PmSOD1 gene of the protistan parasite <i>Perkinsus marinus</i> complements the <i>sod2Delta</i> mutant of <i>Saccharomyces cerevisiae</i> , and directs an iron superoxide dismutase to mitochondria. <i>Mol Biochem Parasitol.</i> 2003. 126:81-92
M. E. Bortoni, V. S. Terra, J. Hinds, P. W. Andrew, H. Yesilkaya. The pneumococcal response to oxidative stress includes a role for Rgg. <i>Microbiology.</i> 2009. 155:4123-34
D. Schafer, T. T. Lam, T. Geiger, M. Mainiero, S. Engelmann, M. Hussain, A. Bosserhoff, M. Frosch, M. Bischoff, C. Wolz, J. Reidl, B. Sinha. A point mutation in the sensor histidine kinase SaeS of <i>Staphylococcus aureus</i> strain Newman alters the response to biocide exposure. <i>J Bacteriol.</i> 2009. 191:7306-14
I. L. Jung, I. G. Kim. Polyamines reduce paraquat-induced <i>soxS</i> and its regulon expression in <i>Escherichia coli</i> . <i>Cell Biol Toxicol.</i> 2003. 19:29-41
J. Wu, Z. Jiang, M. Liu, X. Gong, S. Wu, C. M. Burns, Z. Li. Polynucleotide phosphorylase protects <i>Escherichia coli</i> against oxidative stress. <i>Biochemistry.</i> 2009. 48:2012-20
J. H. Chou, J. T. Greenberg, B. Dimple. Posttranscriptional repression of <i>Escherichia coli</i> OmpF protein in response to redox stress: positive control of the <i>micF</i> antisense RNA by the <i>soxRS</i> locus. <i>J Bacteriol.</i> 1993. 175:1026-31

Reactive oxygen species (ROS) from a variety of sources are often encountered by invading plant pathogens during the in
The cytotoxic effects of reactive oxygen species are largely mediated by iron. Hydrogen peroxide reacts with iron to form
<i>Pseudomonas chlororaphis</i> GP72 is an important plant growth-promoting rhizobacteria (PGPR) with a wide-spectrum ant
Extracts of <i>E. coli</i> contain at least three easily separable NAD(P)H:paraquat diaphorases. One of these is identified as thio
We report the first example of a gene, <i>hmp</i> , encoding a soluble flavohemoglobin in <i>Escherichia coli</i> K-12, which is up-reg
<i>Streptococcus mutans</i> , a key etiological agent of human dental caries, lives almost exclusively on the tooth surface in pla
Thioredoxin-like proteins of the TlpA/ResE/CcmG subfamily are known to face the periplasm in gram-negative bacteria. U
High coverage, whole genome shotgun (WGS) sequencing of 57 geographically- and genetically-diverse isolates of <i>Strept</i>
A method to grow the halophilic archaeon <i>Haloferax volcanii</i> in microtiter plates has been optimized and now allows the
The lipid peroxidation product 4-hydroxynon-2-enal (4-HNE) is a strong electrophile that forms covalent adducts with pro
Expression of plant metallothionein genes has been reported in a variety of senescing tissues, such as leaves and stems, t
Insertional mutagenesis was used in <i>Chlamydomonas reinhardtii</i> to isolate original mutants hypersensitive to multiple dr
The facultative intracellular oyster parasite, <i>Perkinsus marinus</i> , taxonomically related to both dinoflagellates and apicom
<i>Streptococcus pneumoniae</i> resides in the oxygen-rich environment of the upper respiratory tract, and therefore the abili
<i>Staphylococcus aureus</i> reacts to changing environmental conditions such as heat, pH, and chemicals through global regu
Polyamines, ubiquitous polycationic compounds, are involved in many cellular responses and relieve paraquat-induced c
<i>Escherichia coli</i> polynucleotide phosphorylase (PNPase) primarily functions in RNA degradation. It is an exoribonuclease
The <i>soxRS</i> regulon is a cornerstone of the adaptive defense systems of <i>Escherichia coli</i> against oxidative stress. Unexpect

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant

Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1



2362
2363
2364
2365
2366
2367
2368
2369
2370
2371
2372
2373
2374
2375
2376
2377

T. Nunoshiba, B. Demple. Potent intracellular oxidative stress exerted by the carcinogen 4-nitroquinoline-N-oxide. <i>Cancer Res.</i> 1993. 53:3250-2
S. Dong, H. Hu, Y. Wang, Z. Xu, Y. Zha, X. Cai, L. Peng, S. Feng. A pqr2 mutant encodes a defective polyamine transporter and is negatively affected by ABA for paraquat resistance in <i>Arabidopsis thaliana</i> . <i>J Plant Res.</i> 2016. 129:899-907
W. Rungrasamee, K. C. Ryan, M. J. Maroney, P. J. Pomposiello. The PqrR transcriptional repressor of <i>Pseudomonas aeruginosa</i> transduces redox signals via an iron-containing prosthetic group. <i>J Bacteriol.</i> 2009. 191:6709-21
M. Baier, K. J. Dietz. Primary structure and expression of plant homologues of animal and fungal thioredoxin-dependent peroxide reductases and bacterial alkyl hydroperoxide reductases. <i>Plant Mol Biol.</i> 1996. 31:553-64
C. Sanchez, A. J. Gates, G. E. Meakin, T. Uchiumi, L. Girard, D. J. Richardson, E. J. Bedmar, M. J. Delgado. Production of nitric oxide and nitrosylhemoglobin complexes in soybean nodules in response to flooding. <i>Mol Plant Microbe Interact.</i> 2010. 23:702-11
D. R. Agius, W. H. Bannister, R. Balzan. Prokaryotic iron superoxide dismutase replaces cytosolic copper, zinc superoxide dismutase in protecting yeast cells against oxidative stress. <i>Biochem Mol Biol Int.</i> 1998. 44:41-9
H. S. Amash, O. R. Brown, V. A. Padron. Protection by selective amino acid solutions against doxorubicin induced growth inhibition of <i>Escherichia coli</i> . <i>Gen Pharmacol.</i> 1995. 26:983-7
Z. Fang, P. C. Dos Santos. Protective role of bacillithiol in superoxide stress and Fe-S metabolism in <i>Bacillus subtilis</i> . <i>Microbiologyopen.</i> 2015. 4:616-31
C. S. Hwang, Y. U. Baek, H. S. Yim, S. O. Kang. Protective roles of mitochondrial manganese-containing superoxide dismutase against various stresses in <i>Candida albicans</i> . <i>Yeast.</i> 2003. 20:929-41
T. Tam le, H. Antelmann, C. Eymann, D. Albrecht, J. Bernhardt, M. Hecker. Proteome signatures for stress and starvation in <i>Bacillus subtilis</i> as revealed by a 2-D gel image color coding approach. <i>Proteomics.</i> 2006. 6:4565-85
V. Pernikarova, V. Sedlacek, D. Potesil, I. Prochazkova, Z. Zdrahal, P. Bouchal, I. Kucera. Proteome-wide dataset generated by iTRAQ-3DLCMS/MS technique for studying the role of FerB protein in oxidative stress in <i>Paracoccus denitrificans</i> . <i>Data Brief.</i> 2015. 4:390-4
C. Wolf, F. Hochgrafe, H. Kusch, D. Albrecht, M. Hecker, S. Engelmann. Proteomic analysis of antioxidant strategies of <i>Staphylococcus aureus</i> : diverse responses to different oxidants. <i>Proteomics.</i> 2008. 8:3139-53
T. Takac, O. Samajova, P. Vadovic, T. Pechan, P. Kosutova, M. Ovecká, A. Husickova, G. Komis, J. Samaj. Proteomic and biochemical analyses show a functional network of proteins involved in antioxidant defense of the <i>Arabidopsis anp2anp3</i> double mutant. <i>J Proteome Res.</i> 2014. 13:5347-61
V. Pernikarova, V. Sedlacek, D. Potesil, I. Prochazkova, Z. Zdrahal, P. Bouchal, I. Kucera. Proteomic responses to a methyl viologen-induced oxidative stress in the wild type and FerB mutant strains of <i>Paracoccus denitrificans</i> . <i>J Proteomics.</i> 2015. 125:68-75
N. J. Hare, N. E. Scott, E. H. Shin, A. M. Connolly, M. R. Larsen, G. Palmisano, S. J. Cordwell. Proteomics of the oxidative stress response induced by hydrogen peroxide and paraquat reveals a novel AhpC-like protein in <i>Pseudomonas aeruginosa</i> . <i>Proteomics.</i> 2011. 11:3056-69
L. N. Nefedova, S. Fantin lu, V. V. Zinchenko, M. M. Babykin. [The prqA and mvrA genes encoding carrier proteins control resistance to methyl viologen in the cyanobacterium <i>Synechocystis</i> sp. PCC6803]. <i>Genetika.</i> 2003. 39:336-40

Oxidative stress exerted by superoxide-generating (redox-cycling) agents such as paraquat triggers the soxRS regulon of <i>E. coli</i>
Despite the paraquat-resistant mutants that have been reported in plants, this study identified a novel <i>A. thaliana</i> mutant that is resistant to paraquat
Inducible defenses against oxidative stress are coordinated by redox-sensitive transcription factors that transduce oxidative stress signals
Higher plants express genes encoding peroxiredoxins of the two-cysteine type. This is concluded from the isolation of cDNA clones from Arabidopsis thaliana
Nitric oxide (NO) has gained interest as a major signaling molecule during plant development and in response to environmental stresses
The iron superoxide dismutase (FeSOD) gene of <i>Escherichia coli</i> was cloned in <i>Saccharomyces cerevisiae</i> cells deficient in iron superoxide dismutase
1. Incubation of <i>Escherichia coli</i> with 0.7 mM doxorubicin in MBS-glucose medium resulted in complete growth inhibition
Glutathione (GSH) serves as the prime thiol in most organisms as its depletion increases antibiotic and metal toxicity, implying a protective role for GSH
<i>Candida albicans</i> contains copper- and zinc-containing superoxide dismutase but also two manganese-containing superoxide dismutases
In this paper we have defined proteome signatures of <i>Bacillus subtilis</i> in response to heat, salt, peroxide, and superoxide
3DLC protein- and peptide-fractionation technique combined with iTRAQ-peptide labeling and Orbitrap mass spectrometry for proteomic analysis
The high resolution 2-D protein gel electrophoresis technique combined with MALDI-TOF MS and a recently developed fluorescence-activated cell sorting (FACS) method
Disentanglement of functional complexity associated with plant mitogen-activated protein kinase (MAPK) signaling has been achieved
FerB is a cytoplasmic flavoprotein from the soil bacterium <i>Paracoccus denitrificans</i> with a putative role in defense against oxidative stress
<i>Pseudomonas aeruginosa</i> is a ubiquitous pathogen most typically associated with wound infections, but also the main cause of nosocomial infections
Derivatives with insertional inactivation of <i>prqA</i> and <i>mvrA</i> genes were obtained and studied in the <i>Synechocystis</i> sp. PCC 6803

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant